



## Agromet Advisory Bulletin

Date : 28-09-2021

Weather Forecast of **KATOL** Block in **NAGPUR**(Maharashtra) Issued On :2021-09-28(Valid Till 08:30 IST of the next 5 days)

Parameter	2021-09-29	2021-09-30	2021-10-01	2021-10-02	2021-10-03
Rainfall	7.6	2.4	4.7	6.0	7.1
Tmax(°C)	31.6	31.8	32.1	32.3	32.5
Tmin(°C)	24.6	24.4	24.9	24.7	24.8
RH-I(%)	94	92	90	88	87
RH-II(%)	77	75	73	72	71
Wind Speed(kmph)	13.0	10.0	8.0	7.0	6.0
Wind Direction(Degree)	169	169	248	23	211
Cloud Cover(Octa)	7	8	7	7	8

### Weather Summary/Alert:

In Katol block, as per the block level value added forecast given by, IMD, RMC, Nagpur, sky will be partially to mostly cloudy and very light to light rain occur during next five days during 29th September, 2021 to 03rd, October, 2021 and thunderstorm accompanied with lightning and gusty winds (speed reaching 30-40 kmph) very likely to occur 29th September, 2021 is forecasted.

### General Advisory:

The spraying of agrochemicals in field crop, vegetables and fruit orchards should be carry out by judging the local weather condition within next 2 to 3 day. Carry out harvesting of matured soybean and others crop and harvested should be stored on elevated field and if threshing is not possible and cover the harvested produce with tarpaulin and allow for sun drying by considering the local weather condition.

### SMS Advisory:

Considering the gusty wind forecast, support the vegetables and orchards with the help of bamboo.

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
SOYABEAN	Soybean harvesting and threshing management: Drain out excess water from the Soybean field if rain water stagnated. Farmers who have grown early maturing soybean varieties like JS 95-60, JS 20-34, JS 93-05, NRC 130, NRC 138 etc. are advised to harvest the crop immediately after attaining the physiological maturity (change in pod colour) and the harvested crop should be stored on elevated field and if threshing is not possible and cover the harvested produce with tarpaulin. Because of continued rain during the maturity stage, the soybean crop is likely to be affected in terms of its quality parameters including viability as well as risk of vivipary in the matured pods. Therefore, farmers are advised to harvest their crop at the right time (Physiological maturity indicating change in pod colour). Farmers are suggested to allow the harvested crop for sun-drying for 2-3 days. It is also advised to collect the harvested crop and keep it in safe place, covering with Tarpaulin and thresh the same as per the convenience.

Crop(Varieties)	Crop Specific Advisory
SORGHUM (JOWAR/GREAT MILLET)	Rabi sorghum land preparation, seed treatment, nutrient and sowing management: Medium to deep / heavy, high water holding capacity and well-drained soil should be selected for rabi sorghum sowing. After harvesting black gram and green gram crops in kharif season, carry out preparation of the land by giving 3 to 4 harrowing. Apply 10 to 15 carts of well decomposed FYM or compost manure before the last harrowing. The land should be prepared for sowing by removing weeds and stubbles. Use 10 kg certified seed per hectare for sowing of rabi sorghum. For sowing, the spacing between two rows should be 45 cm and between two plants 15 cm. For expected yield of irrigated rabi sorghum, the spacing between two rows should be 45 cm and between two plants 12 cm. For sowing use C.S.H. – 15 R hybrid variety while improved / pure variety PKV Kranti (AKSV 13 R), Parbhani Moti, CSV-18 (Rabbi Irrigated Variety), CSV- 12, CSV- 29, Phule Vasudha, Phule Suchitra, Phule Revati, Parbhani Super Moti and local varieties are Maldandi: 35-1 and Ringani should be used. Sorghum seeds should be treated with 25 g of Azotobacter, 20 g of Phosphorus Solubilizing Bacteria (PSB) and 4 g of Trichoderma viride per kg of seed. For dryland rabi sorghum, apply 50 kg Nitrogen (N), 25 kg Phosphorous (P) and 25 kg Potassium (K) per hectare at the time of sowing. Sowing should be done across the slope. Considering the available soil moisture and after arrival of Wafsa condition, sowing of rabi sorghum should be done between 25th September to 15th October.
COTTON	Cotton disease management: Drain out excess water from the cottonfield if rain water stagnated. As precautionary measure, prophylactic spraying should be done in clear weather after current spell of rain with Copper oxychloride 50 WP @25 g+ [Streptomycin sulphate IP 90% w/w+ Tetracycline hydrochloride IP 10% w/w] @ 1 g in 10 litres of water is suggested during early boll developmental stages for the management of internal boll rot and bacterial leaf blight of cotton. If symptoms of parawilting observed in cotton, farmers are advised to undertake drenching of carbendazim 50 WP @25g + Urea @100g in 10 litres of water after current spell of rain. Cotton sucking pest management: Infestation of sucking pests (Thrips and Jassid) should be monitored during squaring, flowering and early boll development stages and for their management spray Thiamethoxam 25 WG @ 2 g or Dinotefuran 20 SG @ 3 g or Diafenthiuron 50 WP @ 12 g per 10 litres of water (Knapsack or battery -operated sprayer). If thrips infestation is prominent (vector of TSV disease) then spray Thiamethoxam 25 WG @2 g or Spinetoram 11.7 SC @8.5 ml or Dinotefuran 20 SG @3 g/10 litres of water after current spell of rain. Cotton pink boll worm management: In order to monitor the pink bollworm moth's activity in cotton, used two pheromone trap per acre with active ingredient of pectinolure or gossypure. Regular observations of moths trapped in pheromone traps should be recorded. As soon as it is observed that the moths have crossed economic threshold level (8 moths per trap per night for three consecutive nights), the recommended chemical pesticides should be applied. Infected rosette flower should be removed and destroyed immediately (Economic Threshold Level– 10 % infested inflorescence flowers). After formation of green bolls, 20 bolls per acre (1 boll per tree) should be monitored randomly every week. (Economic Threshold Level- 10 % affected bolls). Depending on availability, Trichogramma bacterium, a parasitic friendly insect that feeds on eggs, should be release three times at an interval of 15 days during the square formation and flowering stage at the rate of 60,000 per acre. Chemical pesticides should not be used at least one week before and one week after use of biological agents for pest control. Apply neem-based insecticides @50 ml or Chlorpyrifos 20% EC @25 ml or Quinalphos 20 AF @20 ml or Thiodicarb 75 WP @20 g or Profenophos 50 EC @30 ml or Indoxacarb 14.5 SC @5 ml or Emamectin benzoate 5 SG @5 g per 10 litres of water wherever pink bollworm crosses ETL after current spell of rain. Cotton nutrient management: Spray 2% urea at flowering stage and 2% DAP at boll development stage of cotton after current spell of rain. Apply second split of 90 kg urea per ha for irrigated hybrid Bt after arrival of Wafsa condition. Cotton square & flower dropping and excess vegetative growth management: It is suggested to spray alpha-NAA 4.5 SL @ 5 ml /10 litres of water to avoid natural sheddi
PIGEON PEA (RED GRAM/ARHAR)	Drain out excess water from the Pigeon pea field if rain water stagnated. Due to continued rainy weather/waterlogging, wilting symptoms may be observed in pigeon peacrop, for its management undertake drenching with copper oxychloride 50%WP@ 25 g in 10 litres of water during clear weather. Install pheromone traps @ 5/ha for monitoring of Helicoverpa. Monitor for presence of eggs & small larvae.

Crop(Varieties)	Crop Specific Advisory
CITRUS	<p>Management of fruit dropping in Nagpur mandarin and sweet orange in rainy season: Excess water should be drained immediately if water is stagnant in orange and sweet orange orchards. Remove water shoots from the plant and all shoots below the bud union of the graft. Keep the orchards weed free. Foliar spray application of GA3 1.5 g + KNO<sub>3</sub> 1.5 kg in 100 litres of water has already be done in first week of September, Foliar spray application of 2,4-D or NAA 1.5 g + monopotassium phosphate (00:52:34) 1.5 kg/100 litres of water in third week of September. Application of 50 g FeSO<sub>4</sub> and 50g ZnSO<sub>4</sub>/plant mixed with 5 kg FYM or vermicompost along with remaining dose of recommended fertilizers should be applied with in the basin of these trees.Application of tetracycline hydrochloride 600 ppm (60 g / 100 litres of water) followed by Streptocycline or K-cycline 150 ppm (1.5g / 10 litres of water) after 21 days in case of severely greening-infected trees (showing typical greening fruit symptoms with reddening of fruit button tissue). Foliar spraying of fosetyl Al 2.5 g per litre of water (twice during August and September at 40 days interval) should be undertaken for Phytophthora brown rot of fruits. Soil drenching + foliar spraying of mefenoxam MZ @ 2.5 g/lit (5-10 litres depending on the tree size) per plant or, foliar spraying of fosetyl Al /Aliette @ 2.5 g/lit (twice at 40 days interval) for management of root rot. All the fruits should be removed followed by soil drenching with mefenoxam MZ (2.5 g/ L) + carbendazim (1 g/ L) solution for root rot affected trees showing severe wilting -type symptoms. For the fruit drop associated with Colletotrichum and Botryodiplodia, the affected trees should be sprayed (twice) with carbendazim or thiophanate methyl at the concentration of 0.1 % (i.e. 1 g /lt). The second spray should be repeated after 10 days of first spraying. Spraying of (azoxystrobin + difenoconazole) combination fungicide @ 0.5 ml/lit can be taken up as an alternative. Installation of methyl eugenol traps at the rate of 20-25 traps per hectare for mass trapping of fruit flies and for effective mass trapping change the lure at every 15-20 days interval. Foliar application of neem oil 1% or petroleum spray oil (Horticulture Mineral Oil) @ 2% at 10-15 days interval during colour break stage, till the harvest for management of fruit sucking moth. Collection of fallen fruits and their destruction at regular intervals would prevent the development of puparia and thus reduce the fruit fly population for next year.All spraying operation should be done after current spell rain and fertilizer operations should be done after arrival of Wafsa condition.</p>
RICE	<p>Paddy plant hoppers management: For management of plant hoppers, although rice crop is prone to plant hoppers, Azadiractin 0.15% (1500 ppm) @ 30 ml or Metarhizium anisopliae as a bio-insecticide @ 2.5 kg/ha. or Imidacloprid 17.8 SL.@ 2.2 ml. or Fipronil 5 SC@ 20 ml. or Flonicamid 50 WG @ 3.0 gm Mix in 10 liters of water and spray. Paddy nutrient management:Apply 100 kg Nitrogen, 50 kg Phosphorus and 50 kg Potash per hectare. Apply whole phosphorus and potash and half the amount of nitrogen in the mud (Puddled field) and the remaining half nitrogen in two equal instalments (usually tillering stage which comes at 30 days and panicle initiation stage which comes about 60 days). Paddy blast disease management: For management of blast of paddy spray Hexaconazole 5% EC @ 20 ml. or Isoprothiolane 40% EC @ 15 ml or Tebuconazole 25.9% EC @ 15 ml per 10 liters of water during clear weather condition. Paddy bacterial leaf blight disease management: For management of Bacterial leaf blight spray Streptocycline 1 to 1.5 gm per 10 liters of water. Paddy sheath blight disease management: For management of Sheath blight spray Tebuconazole 25.9% EC @ 15 ml or Propiconazole 25% EC @ 10 ml or Hexaconazole 5 EC @ 20 ml by mixing in 10 liters of water. Paddy sheath rot disease management: For management of sheath rot, Spray Hexaconazole 75 WP @ 1.33gm by mixing in 10 liters of water. Paddy grain discoloration management: For management of grain discoloration, While the crop is in flowering stage, spray any one of the following fungicides in the afternoon. Hexaconazole 4% + Zyneb 68% W.P.@ 20 to 25 gm or Tebuconazole 50% + Trifloxystrobin 25% WG. @ 4 gm per 10 liters of water. All spraying operation should be done after current spell rain.</p>

#### Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
CHILLI	<p>In chilli crop undertake spray of Difenoconazole 25 EC@ 5 ml OR Azoxystrobin 23% SC@ 10 ml in 10 litres of water for control of dieback and fruit rot disease. Undertake spray after current spell of rains.</p>

#### Live Stock Specific Advisory:

<b>Live Stock(Varieties)</b>	<b>Live Stock Specific Advisory</b>
GOAT	Due to rainy season, in goat helminthic parasites and coccidia protozoa infection is on rising trend. Therefore, it is recommended to administer anthelmintic and anticoccidial medication.

**Poultry Specific Advisory:**

<b>Poultry(Varieties)</b>	<b>Poultry Specific Advisory</b>
CHICKEN	Tulsi leaves powder @ 2 gram per days upto 15 days given to poultry to increase immunization power.

**Others (Soil / Land Preparation) Specific Advisory:**

<b>Others (Soil / Land Preparation) (Varieties)</b>	<b>Others (Soil / Land Preparation) Specific Advisory</b>
GENERAL ADVICE	Farmers should take care of thunder and lightning in the sky: If you are working in the field, take shelter immediately near the field. After taking shelter in a safe place in the field, keep dry wood, plastic, gonapath, dry mulch under the feet. Sit with both feet together and both hands on your knees. Make sure that no part of your body touches the ground except your feet. Individuals working in ponds, such as places where there is moisture in the soil or water sources, should go to a safe and dry place immediately. If there are tall trees nearby, take shelter at a distance twice the height of that tree. A pucca house is the safest place to avoid lightning. Farmers should plant trees as low as possible around their houses and livestock sheds. Moist, swampy places and water sources (wells, lakes, rivers, farm ponds etc.) should be avoided as much as possible. If traveling in a four-wheeler, stop in the vehicle. Farmers should take care that they do not have any metal tools. When working in the field, do not let more people work together at the same time. Care should be taken to keep a distance of at least 15-20 feet between two persons. Do not use an umbrella with a metal rod. Stay away from other electric tools made by farmers as well as metal ones.
GENERAL ADVICE	It should be noted that the validity of the dates mentioned in the Weather Based AdvisoryBulletins will be valid from 08:30 AM on the previous day to 08:30 AM on the said date.